

## The Hamiltonian formulation of classical mechanics

Recall the Euler-Lagrange equation in our previous article. We had:

$$\frac{\partial L}{\partial q^i} - \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}^i} \right) = 0 \quad (1)$$

If we define the “conjugate momentum” of  $q^i$  as follows:

$$p_i \equiv \frac{\partial L}{\partial \dot{q}^i} \quad (2)$$

we can write the Euler-Lagrange equation as:

$$\dot{p}_i = \frac{\partial L}{\partial q^i} \quad (3)$$

This allows us to express the small variation of Lagrangian as follows:

$$\begin{aligned} \delta L &= \sum_i \left( \frac{\partial L}{\partial q^i} \delta q^i + \frac{\partial L}{\partial \dot{q}^i} \delta \dot{q}^i \right) \\ &= \sum_i \dot{p}_i \delta q^i + p_i \delta \dot{q}^i \end{aligned} \quad (4)$$

Stepping further, we get:

$$\begin{aligned} \delta L &= \sum_i \dot{p}_i \delta q^i + \delta(p_i \dot{q}^i) - \dot{q}^i \delta p_i \\ \delta \left( \sum_i p_i \dot{q}^i - L \right) &= \sum_i -\dot{p}_i \delta q^i + \dot{q}^i \delta p_i \end{aligned} \quad (5)$$

We now define

$$H \equiv \sum_i p_i \dot{q}^i - L \quad (6)$$

and call it “Hamiltonian.” It turns out that this coincides with energy, as we will see in the next article. At this point, I want to caution the readers that  $H$  must be re-expressed solely in terms of  $p_i$ s and  $q^i$ s by eliminating  $\dot{q}^i$ . In other words,  $\dot{q}^i$  must be absent in  $H$ .

From (5) and (6), we have:

$$\delta H = \sum_i -\dot{p}_i \delta q^i + \dot{q}^i \delta p_i \quad (7)$$

which implies

$$\frac{\partial H}{\partial q^i} = -\dot{p}_i, \quad \frac{\partial H}{\partial p_i} = \dot{q}^i \quad (8)$$

These are called Hamilton's equations.

Let me give you an example. In 3-dimensional Cartesian coordinate, we have:

$$L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - V(x, y, z) \quad (9)$$

Here, we will set:

$$q^1 = x, \quad q^2 = y, \quad q^3 = z \quad (10)$$

Then, from (2), we have:

$$p_1 = m\dot{q}^1, \quad p_2 = m\dot{q}^2, \quad p_3 = m\dot{q}^3 \quad (11)$$

In other words:

$$p_i = m\dot{q}^i \quad (12)$$

Now, we can re-express Lagrangian in terms of  $p$ s and  $q$ s, instead of  $\dot{q}$ s and  $q$ s as follows:

$$L = \sum_i \frac{(p_i)^2}{2m} - V(q_i) \quad (13)$$

Plugging this into (6), we get:

$$\begin{aligned} H &= \sum_i p_i \left( \frac{p_i}{m} \right) - L \\ &= \sum_i \frac{(p_i)^2}{2m} + V(q^i) \end{aligned} \quad (14)$$

Then, Hamilton's equations become:

$$\frac{\partial H}{\partial p_i} = \frac{p_i}{m} = \dot{q}^i \quad (15)$$

$$p_i = m\dot{q}_i \quad (16)$$

$$\frac{\partial H}{\partial q^i} = \frac{\partial V}{\partial q^i} = -\dot{p}_i = -m\ddot{q}^i \quad (17)$$

$$m\ddot{q}^i = -\frac{\partial V}{\partial q^i} \quad (18)$$

So, we recover Newton's equations. Of course, Hamilton's equations are always valid for general cases as well.

Let me conclude this article with a comment. The process in which we obtained Hamiltonian from Lagrangian may seem like a magic but it is

just an example of what is called “Legendre transformation.” The Legendre transformation of function  $f(x, y)$  is given by

$$g(p, y) = f - px \tag{19}$$

where  $p = \frac{\partial f}{\partial x}$ . In our case we had  $x = \dot{q}$ ,  $y = q$ ,  $f = L$ , and  $g = -H$ . Besides the Hamiltonian mechanics, Legendre transformation has applications in thermodynamics and quantum field theory.

**Problem 1.** In our earlier article “Central force problem solution in terms of Lagrangian mechanics,” we obtained the equation of motion for the case in which the Lagrangian was given as follows:

$$L = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\theta}^2) - V(r) \tag{20}$$

First, obtain the Hamiltonian in terms of  $r, \theta, \dot{r}$  and  $\dot{\theta}$ , in such a case and check that it coincides with the usual energy. (i.e. the sum of the kinetic energy and the potential energy) Then, re-express the Hamiltonian in terms of  $r, \theta, p_r$  and  $p_\theta$ , and obtain the equation of motion using Hamilton’s equations and check that they coincide with the earlier ones obtained using Lagrangian formulation of classical mechanics.

## Summary

- The conjugate momentum of  $q^i$  is given by

$$p_i \equiv \frac{\partial L}{\partial \dot{q}^i}$$

- The Hamiltonian is then given by

$$H = \sum_i p_i \dot{q}^i - L$$

- Hamilton’s equations are given by

$$\frac{\partial H}{\partial q^i} = -\dot{p}_i, \quad \frac{\partial H}{\partial p_i} = \dot{q}^i$$

- The Hamiltonian corresponds to the energy (i.e. the sum of kinetic energy and potential energy).